

12 **EUROPEAN PATENT APPLICATION**

21 Application number: 84115346.3

22 Date of filing: 13.12.84

51 Int. Cl.⁴: **A 61 K 37/54**
 A 61 K 37/547, C 12 N 9/50
 C 12 N 9/68, C 12 N 9/72
 C 12 N 11/02, C 07 D 213/70

30 Priority: 24.12.83 GB 8334499

43 Date of publication of application:
 25.09.85 Bulletin 85/39

84 Designated Contracting States:
 BE CH DE FR GB IT LI NL SE

71 Applicant: **BEECHAM GROUP PLC**
 Beecham House Great West Road
 Brentford Middlesex TW8 9BD(GB)

72 Inventor: **Smith, Richard Anthony Godwin**
 22 Evesham Road
 Reigate Surrey RH2 9DL(GB)

74 Representative: **Hardman, Carol Pauline et al,**
 Beecham Pharmaceuticals Great Burgh Yew Tree
 Bottom Road
 Epsom Surrey KT18 5XQ(GB)

54 Enzyme-protein conjugate, process for its preparation and pharmaceutical use thereof.

57 A derivative of a fibrinolytic enzyme in which the catalytic site on the enzyme which is responsible for fibrinolytic activity is blocked by human protein attached thereto by way of a reversible linking group.




Fig.1

TITLE MODIFIED

NOVEL DERIVATIVES

This invention relates to enzyme derivatives for use in the treatment of thrombotic diseases.

European Patent No. 0,009,879 discloses derivatives of in vivo fibrinolytic enzymes which are useful therapeutic agents for treating venous thrombosis. The derivatives are characterised by the active catalytic site on the enzymes being blocked by a group which is removable by hydrolysis such that the pseudo first order rate constant for hydrolysis is in the range 10^{-6} sec^{-1} to 10^{-3} sec^{-1} .

It has now been found that certain human proteins may be chemically modified so that they can function as removable blocking groups in the sense used in the above mentioned European Patent, and that these groups, when linked to enzymes, can produce enzyme derivatives with slow physiological clearance rates.

According to the present invention there is provided a derivative of a fibrinolytic enzyme in which the catalytic site on the enzyme which is responsible for fibrinolytic activity is blocked by a human protein attached thereto by way of a reversible linking group.

As used herein the expression "reversible linking group" includes groups which are

- 2 -

removable by hydrolysis at a rate such that the pseudo first order rate constant for hydrolysis is in the range 10^{-6} sec^{-1} to 10^{-3} sec^{-1} in isotonic aqueous media at pH 7.4 at 37°C .

The preferred rate constant is in the range $1 \times 10^{-5} \text{ sec}^{-1}$ to $8 \times 10^{-4} \text{ sec}^{-1}$.

Suitably the catalytic site on the enzyme is blocked by group of structure (I)



in which

P is a human protein modified by treatment with an amino acid side chain specific reagent to include a protein attachment group,

X is an acyl group of formula



in which R is an aromatic or aliphatic residue, A is a bridging group comprising at least one hetero atom selected from oxygen, sulphur and nitrogen, in which the nitrogen is optionally substituted by $\text{C}_1\text{-6}$ alkyl,

B is a linear hydrophilic linking group, linked to the protein attachment group on P.

The term 'fibrinolytic enzyme' is used herein to mean any enzyme which demonstrates in vivo fibrinolytic activity as defined in the above mentioned European Patent, and includes enzymes which are obtainable from mammalian urine, blood or tissues, or from bacteria expressing genes specifying such mammalian enzymes and

plasminogen activator, the extraction of which is described in

Published European Patent Application No. 41766. t-PA is a type of activator known as human extrinsic plasminogen activator. Intrinsic plasminogen activators, such as urokinase and plasmin, are also suitable fibrinolytic enzymes for use in the present invention.

Examples of suitable human proteins are human plasma proteins known to have slow physiological clearance rates such as human serum albumin (and its oligomers), immunoglobulin, human fibrinogen and human plasminogen. Preferably the protein has a molecular weight in the region of 30,000 to 800,000 daltons. Preferred proteins are ones which are known to have significant affinity for components of a thrombotic mass, for example, human plasminogen. Optionally, the protein may itself be a fibrinolytic enzyme as hereinbefore defined such as urokinase or tissue plasminogen activator, or a proenzyme thereof such as prourokinase. When the protein is a fibrinolytic enzyme, the active catalytic site thereof may optionally be blocked as described in European Patent No 0009 879. An example of such a blocked fibrinolytic enzyme is p-aminobenzoyl urokinase.

Examples of derivatives of the invention include

human serum albumin → human tissue plasminogen
activator

human serum albumin → urokinase

urokinase or acylated urokinase → human plasmin

- 4 -

human immunoglobulin G \rightarrow human tissue plasminogen
activator


human plasminogen \rightarrow human tissue plasminogen activator

where \rightarrow represents a reversible linkage to the active
centre of the enzyme.

Examples of suitable groups X include groups derived
from those blocking groups described in European Patent
No. 0,009,879. Preferred groups are optionally
substituted benzoyl groups as described in the above
mentioned European Patent further substituted at the 2
or 4 position by the group A, and optionally
substituted acryloyl groups also described in the above
mentioned European patent and joined to A at the 2 or 3
position thereof.

Suitable groups A are those which provide sufficient
stabilisation of the resulting benzoyl or
acryloyl-enzyme to result in a pseudo first order rate
constant for hydrolysis in the above mentioned range
and preferably in the range 1×10^{-5} to $8 \times 10^{-4} \text{ sec}^{-1}$.

Examples of A are:



The protein attachment group is a functionality derived by modification of the protein with a reagent specific for one or more amino acid side chains, and which contains a group capable of reacting with a group B.


Examples of groups B are substituted C₂-C₁₀ alkanes such as 6-aminohexyl, or linear polymers such as polyethylene glycol, polypropylene glycol, poly-glycine, poly-alanine or poly-sarcosine. The linear group B may optionally contain a cleavable section to facilitate analysis of the derivative or to react with the protein attachment group such as those derived from a 3-thio propionyl or 2-thio acetyl derivative of the ω-amino alkane or polymer function. An example of a cleavable section is a disulphide bond. Preferably the disulphide bond is derived from reaction of P with the linear group B and thus is generated at the linkage of B with P. Alternatively the cleavable section may comprise an α, β dihydroxy function.

Suitably B is a group; - S(CH₂)₂CONH - ,
-S-(CH₂)₂CONH(CH₂)₆- or -S-(CH₂)₂CONH(CH₂)₂-

As an example, the generation of a free thiol function by reaction of the protein with 2-iminothiolane or N-acetyl homocysteine thiolactone will permit coupling of the protein attachment group with a thiol-reactive B structure. Alternatively, the protein attachment group can contain a thiol-reactive entity such as the 6-maleimidoethyl group or a 2-pyridyl-dithio group which can react with a free thiol in B. Preferably, the protein attachment group is derived from protein modifying agents such as 2-iminothiolane that react

The derivatives of the present invention may be prepared by reacting together a human protein optionally modified to include a protein attachment group, a fibrinolytic enzyme and a linking agent having a moiety capable of reacting with the catalytic site of the enzyme and a moiety capable of reacting with a protein amino group or protein attachment group to form a reversible linking group as hereinbefore defined.

In particular the derivatives may be prepared by treating a human protein, which has been optionally modified by treatment with an amino acid side chain specific reagent to include a protein attachment group, with an acylating agent of formula (II)



in which B, A and X are as defined with reference to formula (I),

W represents a group capable of reacting directly with the amino acid side chain of a protein or, when the protein includes a protein attachment group, W represents a group capable of reacting with the attachment group, Z represents a counter anion, preferably halide or p-toluenesulphonate, and each of R_1 to R_4 represents hydrogen or an electron withdrawing moiety which increases the reactivity of an amidinophenyl ester;

and reacting the acylated protein produced thereby with a fibrinolytic enzyme.

The acylating agents of formula (II) are novel and as such form part of the invention.

When W represents a group capable of reacting directly with the amino acid side chain of a protein, it is preferably an N-succinimidyl group. When W represents a group capable of reacting with a protein attachment group, it is preferably a pyridylthio group. Optionally W may be a photoactivated group such as 2-nitro-4-azido phenyl.

Preferably, each of R¹ to R⁴ represents hydrogen or halogen.

The derivatives of the present invention may also be prepared by treating a human protein, which has been modified by treatment with an amino acid side chain specific reagent to include a protein attachment group, with a fibrinolytic enzyme which itself has been modified by treatment with an acylating agent of formula (II) in which W is a group capable of reacting with the protein attachment group on the protein.

Preferably, W is a pyridylthio group.

In the above processes, modification of the protein to introduce a protein attachment group is preferably carried out in aqueous buffered media at a pH between 3.0 and 9.0 depending on the reagent used. For a preferred reagent, 2-iminothiolane, the pH is preferably 6.5-8.5. The concentration of protein is preferably high (> 10mg/ml) and the modifying reagent is used in a moderate (1.1- to 5-fold) molar excess, depending on the reactivity of the reagent. The

of modification of the protein may be determined by assaying the protein attachment groups introduced.

- 8 -

Such assays may be standard protein chemical techniques such as titration with 5,5'-dithiobis-(2-nitrobenzoic acid). Preferably, 0.5-2.0 moles of protein attachment group will be introduced on average per mole of protein. The modified protein may be separated from excess modifying agents by standard techniques such as dialysis, ultrafiltration, gel filtration and solvent or salt precipitation. It is generally desirable to react the modified protein with the acylating agent or the acylated fibrinolytic enzyme as soon as possible, but in certain cases, the intermediate material may be stored in frozen solution or lyophilised.

The modified protein prepared as described above may be reacted with the acylating agent of formula (II) under conditions similar to those used for the initial introduction of the protein attachment group but with the following qualifications:

- (a) To avoid hydrolysis of the amidinophenyl ester, the preferred pH range is 7.0 to 8.0 and non-nucleophilic buffers should be used.
- (b) The preferred temperature range is 0°C-30°C and the duration of reaction up to 6 hours.
- (c) The molar ratio of acylating agent to protein attachment group is preferably in the range 1 to 10.
- (d) The reaction may, optionally, be monitored by observing the release of a derivative of group W

(5,5'-dithiobis(2-nitrobenzoate))

- (e) Because of the reactivity of the product (a proteinaceous acylating agent), it is desirable to separate the product from the excess acylating agent of formula (II) as quickly as possible. For this purpose, high performance size exclusion chromatography or diafiltration may be used.
- (f) The product should preferably be reacted with the fibrinolytic enzyme immediately but may be stored in frozen solution (not lyophilised) below -40°C for short periods.

The treatment of an unmodified protein with an acylating agent of formula (II) is generally performed under conditions similar to those used for the introduction of a protein attachment group. However, the reactivity of this type of reagent requires that precautions similar to those noted in paragraphs (a) to (f) above should be exercised.

Where the modified protein itself comprises a fibrinolytic enzyme whose active catalytic site is blocked, the blocking may be carried out as described in European Patent No 000 9879 either before or after modification as described above, or after reaction with the fibrinolytic enzyme.

In the process aspect of the invention which utilises a reaction between modified protein and acylated fibrinolytic enzyme, the enzyme may first be reacted with an acylating agent of formula (II) under the conditions described for the introduction of blocking groups in European Published Patent Application No. 0,009,879. Having been freed of excess reagent by the

group under conditions similar to those used in

- 10 -

paragraphs (a) to (d) above. However, it is preferable to conduct the coupling below 100°C (preferably 00-40°C) in order to minimise the hydrolysis of the acylated enzyme. In addition, the modified protein may be used in a large molar excess (up to 10⁴-fold) over the acylated enzyme. The latter conditions also apply to the coupling between an a proteinaceous acylating agent and fibrinolytic enzyme.

The derivatives of this invention may be purified from excess modified protein/protein acylating agent and uncoupled enzyme by a variety of separation techniques. Because the molecular weight of the derivative will usually be significantly greater than that of its components, rapid size fractionation methods such as protein high performance gel chromatography are particularly advantageous. In addition, affinity chromatography utilising an insoluble matrix with affinity for the fibrinolytic enzyme component, may be used, in conjunction with gel filtration methods.

The acylating agents of formula (II) may be prepared by a variety of standard procedures. A preferred general synthetic route is given below.

- (a) Reaction of an acylating agent containing a masked reactive functionality (for example, N-succinimidyl-3-(2-pyridyldithio)propionate-SPDP) with a 2- or 4- substituted benzoic acid derivative containing a nucleophilic function on the substituents (for example: 4-hydrazinobenzoic acid, N-2 (6-aminohexyl) aminobenzoic acid or N-4 (6-aminohexyl) aminobenzoic acid). Preferred

another possible procedure is to react for 1-24 hours.

- b) The intermediate acid from (a) is esterified with a salt of 4-amidinophenol (or a ring-substituted derivative thereof) using dicyclohexylcarbodiimide in a weakly basic solvent as described for simple blocking agents in European Published Patent No. 0,009,879. It is preferable to use a slight molar excess of the amidinophenol (1.5 to 4-fold) to ensure efficient esterification. Optionally, the esterification may be performed in the presence of anhydrous p-toluenesulphonic acid as an acidic catalyst.

The derivatives of this invention are preferably administered as pharmaceutical compositions.

Accordingly, the present invention also provides a pharmaceutical composition comprising the derivative of the invention in combination with a pharmaceutically acceptable carrier.

The compositions according to the invention may be formulated in accordance with routine procedures as pharmaceutical compositions adapted for intravenous administration to human beings.

Typically compositions for intravenous administration are solutions of the sterile derivative in sterile isotonic aqueous buffer. Where necessary the composition may also include a solubilising agent to keep the derivative in solution and a local anaesthetic such as lignocaine to ease pain at the site of injection. Generally, the derivative will be supplied in unit dosage form for example as a dry powder or water-free concentrate in a hermetically sealed

- 12 -

as an indication of the time within which the free, unmodified protein will be liberated. Where the derivative is to be administered by infusion, the derivative will be dispensed with an infusion bottle containing sterile pharmaceutical 'water for injection'. Where the derivative is to be administered by injection the derivative is dispensed with an ampoule of sterile water for injection. The injectable or infusable composition will be made up by mixing the ingredients prior to administration.

The quantity of material administered will depend upon the amount of fibrinolysis required and the speed with which it is required, the seriousness of the thromboembolic condition and position and size of the clot. The precise dose to be employed and mode of administration must per force in view of the nature of the complaint be decided according to the circumstances by the physician supervising treatment. However, in general, a patient being treated for a mature thrombus will generally receive a daily dose of from 0.10 to 2.0 mg/kg-1 of body weight either by injection in up to five doses or by infusion. For the treatment of coronary thrombosis a similar dose may be given as a single intravenous bolus.

No toxic effects have been observed with the compounds of the invention within the above described dosage range.

Accordingly, in a further aspect of the invention there is provided a method of treating thrombotic disorders, which comprises administering to the sufferer an effective non-toxic amount of a derivative of the

The following Methods and Examples illustrate the invention.

Methods

(a) Chromogenic substrate assay

Urokinase and t-PA were assayed against the chromogenic substrates (KabiVitrum, Sweden) S-2444 and S-2288, respectively at a substrate concentration of 1 mM in 0.1 M triethanolamine.HCl pH 8.0 at 25°C. An SU is defined as the amount of activity that gives an O.D. increase of 0.001/min in 1ml substrate in a 1 cm pathlength cell.

(b) Rate constant determinations

The pseudo first order rate constant is determined by hydrolysing the acyl-enzyme under physiological conditions, i.e. in isotonic aqueous media at pH 7.4 and at 37°C. At regular intervals aliquots are withdrawn and incubated with a chromogenic substrate and the rate of conversion of the substrate measured as indicated above.

The hydrolysis is followed until such time as the rate of conversion of substrate reaches a maximum. The rate constant k is then calculated by plotting:

$$\log_e (1 - A_t/A_{\max}) \text{ against } t$$

where A_{\max} is the maximum rate at which an aliquot converts substrate and A_t is the rate at which an aliquot converts substrate at time t .

(c) Assay of fibrinolytic activity in the bloodstream
of rats

Male Sprague-Dawley rats (300-400g) were anaesthetized with pentobarbitone sodium (60mg/kg i.p.). One carotid artery was cannulated for collection of blood samples. One femoral vein was cannulated for injection of heparin (50 U/kg) and compound under test.

Approximately 5 min after heparinization, a pre-dose blood sample (0.8ml) was taken and mixed with 0.1 volumes 129 mM trisodium citrate. The compound under test was then injected (1 ml/kg) over 10s. Further blood samples were taken exactly 1, 2, 4, 8, 16, 30 and 60 min later. Heparin treatment (50 U/kg) was repeated after the 30 min sample to maintain cannula patency.

All citrated blood samples were kept on ice until the end of each experiment, then centrifuged at 1700g for 15 min at 4° to obtain plasma. The euglobulin fraction was precipitated by adding 0.1 ml each plasma to 1.82 ml ice-cold 0.011% (v/v) acetic acid in water. After 30 min standing in ice, all tubes were centrifuged at 1700 g for 15 min at 4°. The supernatants were poured away, the inner walls of each tube carefully wiped dry and each precipitate redissolved in 0.4 ml 0.1 M triethanolamine HCl buffer, pH 8.0, containing 0.05% (w/v) sodium azide. Aliquots (20 µl) were then applied to fibrin plates in quadruplicate. Fibrin plates were prepared from 0.4% (w/v) human fibrinogen (Nabi, Grade 1, Flow Laboratories, Scotland) dissolved in 0.029 M barbitone in 125 mM NaCl, pH 7.4, pipetted (9ml) into 10 x 10 cm square plastic dishes (Sterilin) and clotted by rapid mixing with 0.3 ml bovine thrombin (50 NIH units/ml, Parke-Davis, UK). Plates were incubated at 37° for 18-24h usually, but longer if required, and

measured using Vernier calipers. All diameter

each sample were averaged, and this mean converted to fibrinolytic activity by reference to a calibration curve. The latter was obtained by adding known amounts of the compound under test to a stock of plasma pooled from at least ten rats. These standards were processed using the same methods and at the same time as the experimental samples. To construct the calibration curve, diameters (mm) were plotted against \log_{10} concentration of compound. The plasma concentration of compound in each experimental sample was expressed as a percentage of that expected on the basis of the dose given and the assumption of 35 ml plasma/kg body weight for each rat.

(d) Assay of [^{125}I]-labelled conjugates in the bloodstream of rats

Aliquots (100 μl) of citrated blood were mixed with 20 μl 1% NaI and 400 μl 20% trichloroacetic acid (TCA) to precipitate protein-bound ^{125}I . After 30 min on ice, the blood was centrifuged and both supernatant and precipitate counted for ^{125}I -content. The percentage total counts in the precipitate was calculated to give TCA-precipitable counts. For each conjugate, the radiometric clearance pattern was obtained by plotting TCA-precipitable counts (as % of theoretical) in the blood against time.

(e) Assay of fibrinolytic activity in the bloodstream of
guinea pigs

Male Dunkin Hartley guinea pigs (350-450 g) were anaesthetized with urethane (25% w/v solution; 6 ml/kg i.p.). One carotid artery was cannulated for collection of blood samples. One femoral vein was cannulated for injection of heparin (50 U/kg i.v.) and compound under test. Approximately 5 min after heparinization, a pre-dose blood sample (2 ml) was taken and mixed with 0.1 volumes 129 mM trisodium citrate. The compound under test was then injected (1 ml/kg) over 10s. Further blood samples were taken exactly 2, 4, 8, 16, 30, 60 and 90 min later. Heparin treatment (50 U/kg i.v.) was repeated after the 30 min sample to maintain cannula patency. All citrated blood samples were kept on ice until the end of each experiment, then centrifuged at 1700 g for 15 min at 4° to obtain plasma. Each plasma sample was diluted 200-fold in phosphate buffered saline, pH 7.4, containing 0.01% (v/v) Tween 80. Aliquots (30 µl) were then applied to fibrin plates in quadruplicate. Fibrin plates were prepared from 0.4% (w/v) human fibrinogen (Kabi, Grade L, Flow Laboratories, Scotland) dissolved in 0.029 M barbitone in 125 mM NaCl, pH 7.4, pipetted (10 ml) into 10 x 10 cm square plastic dishes (Sterilin) and clotted by rapid mixing with 0.3 ml bovine thrombin (50 NIH units/ml, Parke-Davis, U.K.). Plates

each lysis zone, two diameters perpendicular to each other

were measured using Vernier calipers. All diameters for each sample were averaged, and this mean converted to fibrinolytic activity by reference to a calibration curve. The latter was obtained by adding known amounts of the compound under test to the pre-dose plasma of each animal. These standards were processed using the same methods and at the same time as the experimental samples. To construct the calibration curve, diameters (mm) were plotted against \log_{10} concentration of compound. The plasma concentration of compound in each experimental sample was expressed as a percentage of that expected on the basis of the dose given and the assumption of 50 ml plasma/kg body weight for each guinea pig.

EXAMPLE 1

0155388

(a) 4-[N-2-(3-[2-pyridyl] dithiopropionyl) hydrazino]
benzoic acid

SPDP (100 mg, 0.32 mmol) was dissolved in dry pyridine (1.0 ml) and 4-hydrazino benzoic acid (55 mg 0.32 mmol) added. The mixture was warmed to 56°C for 4h and left at ambient temperature overnight. The product was evaporated to dryness and the resulting yellow oil recrystallised from water (6.0 ml) and EtOH (0.5 ml). A gummy solid was deposited on cooling and this was dissolved in ethanol (1.0 ml), evaporated and dried in vacuo to give a glassy solid.

Yield: 72.6 mg (65%)

NMR (MeOH-D⁴). δ : 8.05, irr d, 1H, Pyridine H6. 7.50, overlapping dd, 4H, Pyridine H3 + 4, + benzoyl A₂B₂ qt. 6.9, Quint, 1H, Pyridine H5. 6.45 d, 2H, benzoyl A₂B₂. 2.6 M. 4H. -CH₂CH₂-

Anal: C₁₃H₂₀N₂O₂ (236.37) requires: C: 66.07, H: 8.53, N: 11.85. Found: C: 66.04, H: 8.16, N: 10.62.

(b) 4-[N-2-(3-[2-pyridyl] dithiopropionyl) hydrazino]
4'-amidinophenyl ester.HCl

The above acid (0.21 mmol) was dissolved in warm (30°C) dry pyridine (1.0 ml) containing 4-amidinophenol.HCl (144 mg, 0.83 mmol) and N,N dicyclohexyl carbodiimide (48 mg, 0.21 mmol). The mixture was left at ambient temperature for 48h and then filtered, washing with dry pyridine. The filtrates were evaporated to dryness and the solid triturated with H₂O (10 ml) and saturated brine (1.0 ml). The solid was recrystallised from EtOH: Et₂O

from EtOH: H₂O 1:2 v/v (1 ml) with addition of brine (0.5 ml) to the filtrate. The yellow solid was dried in vacuo over P₂O₅.

Yield: 52.4 mg (52%). M.P. 140-142°C.

NMR (DMSO - d₆). δ : 10.15, 5, 1H, Amido H. 9.35/9.55 d, 4H, Amidine H, 8.5 irr m. Pyridine H. 7.9/7.5, 7.3/6.8 irr d. Aromatic H, 3.15 d, 2H. CH₂, 2.75 d, 2H, CH₂.

Reduction of the material with 1mM dithiothreitol at pH 7.4 and monitoring of pyridine 2-thione release at 343 nm indicated a purity of about 70%.

Example 2

(a) N-6-(aminohexyl) anthranilic acid

2-chlorobenzoic acid (31.4 g, 0.2 mol), 1.6 diaminoethane (55.7 g, 0.5 mol) anhydrous potassium carbonate (25.6 g), n-pentanol (68 ml) and copper powder (0.3 g) were refluxed for 5h. n-pentanol and 1,6 diaminoethane were distilled off and the residue poured into cold water (1.5 l). The mixture was acidified to pH 7.0 with concentrated HCl and cooled to 4°C. The resulting gum was taken up in hot H₂O: EtOH conc. HCl 8:8:1 v/v (850 ml), cooled, filtered and brought to pH 7.0 with 5 N NaOH. The precipitate was dissolved in H₂O: conc. HCl 1:1 v/v (80 ml), diluted to 300 ml with H₂O and with 5N NaOH. A yellow solid was precipitated at 4°C, filtered and dried in vacuo (5 ml).

Yield: 24.4 g (52%). M.P. 192°C (dec).

NMR (DCl/D₂O). δ : 8.20, M, 1H, Anthraniloyl H₆.

broad envelope. 3H backbone CH₂

(b) N-(6-[3-(2-pyridyl)dithiopropionyl]aminoethyl)
anthranilic acid 4'-amidinophenyl ester. HCl

SPDP (200 mg, 0.64 mmol) was dissolved in dry pyridine (1.0 ml) and added to the above aminoethyl anthranilic acid (151 mg, 0.64 mmol) which rapidly dissolved. The solution was allowed to stand at ambient temperature for 4h and then evaporated under reduced pressure. The resulting oil was dissolved in hot 1 M HCl (5 ml) and the solution neutralised with 5 M NaOH (0.5ml), then allowed to cool overnight at 4°C. A pale yellow solid was precipitated and isolated by decantation of the supernatant and drying of the residue over P₂O₅ in vacuo. The product was dissolved in dry pyridine (1.0 ml) and p-amidinophenol. HCl (221 mg, 1.28 mmol) and anhydrous p-toluenesulphonic acid (100mg) added, followed by N,N dicyclohexylcarbodiimide (158 mg, 0.77 mmol). The mixture was stirred overnight at ambient temperature and filtered, washing with dry pyridine (3.0 ml). Combined filtrates were evaporated under reduced pressure to an oil which was recrystallised twice from 20% v/v ethanol/water.

Yield: 154 mg (30%). M.P. approx. 70°C (glass transition)

NMR (DMSO - d₆, 270 MHz). δ : 9.165/9.436, d, 4H, Exch.D₂O amidine H. 8.056/8.021, dd, 1H, Pyridine H₆. 7.95/7.92 and 7.52/7.51, Qt, 4H. Amidinophenol H. 7.6-7.85, M, 2H, Pyridine H₄ + H₅. 7.48-7.55, M, 2H, Pyridine H₃ + amide NH. 7.21-7.25, M, 1H, anthraniloyl H₄. 7.11-7.14, 1H, anthraniloyl H₂. 6.82-6.86, d, 1H, anthraniloyl H₁. 6.67-6.70, d, 1H, anthraniloyl H₃.

1.32-1.61, 2 multiplets. 8H, -(CH₂)₄-

270

0155388

- 20a -

The NMR spectrum also indicated the presence of approximately 0.5 mol of solvent ethanol and 0.7 mol of p-toluenesulphonate ion per mol of product.

Reduction of the material with dithiothreitol at pH 7.4 and monitoring of the pyridine 2-thione release at 343 nm indicated a purity of about 64%.

EXAMPLE 3

- 21 -

Preparation of substituted amidinophenyl esters of
N-4-{N-2-[3-(2-pyridyldithio)propionyl]aminoethyl}amino
benzoic acid

5 (a) t-butyl 4-aminobenzoate

4-Aminobenzoic acid (5.0 g) was suspended in thionyl
chloride (50 ml) and heated to a gentle reflux. After
two hours the solution went completely clear, and so the
reaction was allowed to cool and the thionyl chloride was
10 removed under reduced pressure, the last traces by azeotrope
with dichloromethane (3 x 50 ml). The resulting acid
chloride was dissolved in more dichloromethane (50 ml) and
a solution of t-butanol (15 ml) in dichloromethane (15 ml)
was added to the stirred solution which was cooled in an
15 ice bath. A solid white precipitate was formed - the
hydrochloride salt of the title compound. This was
isolated by evaporation of the dichloromethane, followed
by suspension of the solid in ethyl acetate (100 ml) and
filtration. The material was suspended in 10% aqueous
20 sodium hydrogencarbonate solution (100 ml) and extracted
into dichloromethane (3 x 100 ml). The organic layer was
dried, filtered and evaporated to leave a pale cream solid
(4.67 g, 66%), m.p. 106-109°C (lit. m.p. 109.5°C)

^1H nmr (CDCl_3 , d^6 DMSO) δ : 7.70 (2H, d, $J = 9\text{Hz}$, aryl-H), 6.55 (2H, d, $J = 9\text{Hz}$, aryl-H), 4.25 (2H, brs, $-\text{NH}_2$), and 1.55 (9H, s, CH_3).

5 Infra Red (Nujol mull) 3420, 3350, 3240, 1690, 1640, 1605, 1515, 1295, 1160, 1120, 850, 770, 700, and 615 cm^{-1} .

(b) t-butyl N-4-(N-2-phthalimidoethyl) aminobenzoate

10 t-Butyl 4-aminobenzoate (1.93 g, 10 mmole) and N-(2-bromoethyl)phthalimide (2.54 g, 10 mmole) were heated together under nitrogen at 100°C for $3\frac{1}{2}\text{h}$. On cooling, the material was dissolved in dichloromethane (125 ml), filtered and evaporated to leave a yellow oil (3.35 g). This was chromatographed (30 g silica/dichloromethane) and the title compound (163 mg, 5%) isolated as a white solid. Recrystallisation from chloroform/petroleum ether ($40-60^\circ\text{C}$)
15 gave a solid m.p. $139-141^\circ\text{C}$.

Found: C 69.70, H 6.14, N 7.71% $\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_4$ requires C 68.84, H 6.05 and N 7.65%

20 ^1H nmr (CDCl_3) δ : 7.6 (5H, m, aryl-H), 6.45 (2H, d, $J = 9\text{Hz}$, aryl-H), 4.5 (1H, br s, N-H), 3.9 (2H, t, $J = 6\text{Hz}$, $\text{CH}_2\text{N}=\text{C}=\text{O}$), 3.4 (2H, m, CH_2NH) and 1.5 (9H, s, CCH_3).

(c) t-butyl N-4-(2-aminoethyl)aminobenzoate

The phthalimide produced above (240 mg) was dissolved in ethanol (10 ml) and hydrazine hydrate (40 μ l) was added. The solution was heated to reflux overnight under nitrogen during which time a white precipitate was formed. Concentrated hydrochloric acid (150 μ l) was added, and the solution cooled and filtered. The ethanol was removed by evaporation to leave an aqueous solution that was refiltered. The solution was basified with 15% sodium hydroxide solution and extracted with ethyl acetate (2 x 25 ml). The organic solution was dried (sodium sulphate), filtered and evaporated to leave a white crystalline compound (131 mg, 85%) - the desired diaminoester, m.p. 117-119°C.

^1H nmr (CDCl_3) δ : 7.75 (2H, d, J = 9Hz, aryl-H), 6.5 (2H, d, J = 9Hz, aryl-H), 4.55 (1H, brs, aryl-NH), 3.25 (4H, brs, CH_2), 1.55 (9H, s, CCH_3), and 1.4 (2H, brs, NH_2)

Infra Red (Nujol) 3370, 1680, 1610, 1540, 1300, 1160, 930, 835, and 775 cm^{-1}

Found C 65.94, H 8.55 and N 11.51%. $\text{C}_{13}\text{H}_{20}\text{N}_2\text{O}_2$ requires C 65.92, H 8.52 and N 11.85%.

(d) t-butyl N-4-{N-2-[3-(2-pyridyldithio)propionyl]
aminoethyl}aminobenzoate

5 t-Butyl N-4-(2-aminoethyl)-aminobenzoate (74 mg) and
N-succinimidyl-3-(2-pyridyldithio)propionate (100 mg)
 were dissolved in dry pyridine (0.5 ml) and stirred
 overnight. The solvent was removed at reduced pressure
 and the residual gel was taken up in ethyl acetate (10 ml)
 and washed with a 10% aqueous sodium hydrogen carbonate
 10 solution (10 ml). The organic layer was dried (sodium
 sulphate), filtered and evaporated to yield an oil (131 mg,
 96%), the title compound.

¹H nmr (CDCl₃) δ: 8.25 (1H, m, aryl-H), 7.4 (6H, m, aryl-
H + NH CO), 6.5 (2H, d, J = 9Hz, aryl-H), 4.65 (1H, brs,
 N-H), 3.4 (4H, m, CH₂NH), 3.05 (2H, t, J = 6Hz, CH₂CO),
 15 2.60 (2H, t, J = 6Hz, CH₂-S-S), and 1.55 (9H, s, CCH₃).

Infra Red (Neat) 3350, 3070, 1660, 1610, 1530, 1420, 1300,
 1160, 1120, 910, 840, 770, 735 cm⁻¹.

(e) N-4-{N-2-[3-(2-pyridyldithio)propionyl]aminoethyl}
aminobenzoic acid, bistrifluoroacetate salt

20 The t-butyl ester (66 mg) was dissolved in trifluoroacetic
 acid (2 ml) and allowed to stand at room temperature for
 reduced pressure and the acid 11 mg
 oil in quantitative yield.

^1H nmr (d^6 acetone/ CDCl_3) δ : 8.6 (1H, m, aryl-H),
 7.75 (5H, m, aryl-H), 6.65 (2H, d, $J = 9\text{Hz}$, aryl-H),
 3.5 (4H, brs, NHCH_2), 3.05 (2H, t, $J = 6\text{Hz}$, COCH_2), and
 2.65 (2H, t, $J = 6\text{Hz}$, CH_2 S).

- 5 (f) 4-amidinophenyl N-4-{N-2-[3-(2-pyridyldithio)
propionyl]aminoethyl}aminobenzoate

The acid bistrifluoroacetate (180 mg) was dissolved in
 pyridine (1 ml). 4-Amidinophenol (54 mg, 1 eq) was added
 followed by dicyclohexyl carbodiimide (64 mg, 1 eq). The
 10 solution went deep orange but this soon dissipated and
 dicyclohexyl urea was rapidly formed. After 5h the
 solution was filtered through a plug of glass wool and
 pyridine was removed by evaporation to leave a brown oil
 (365 mg). Assuming complete esterification, this material
 15 was ~ 45% by weight the desired material and was contaminated
 by pyridinium trifluoroacetate.

Infra Red (Neat) 2400-3300, 1680, 1610, 1490, 1170, 840,
 750 and 710 cm^{-1}

- 20 (g) 2-Chloro-4-amidinophenyl N-4-{N-2-[3-(2-pyridyldithio)
propionyl]aminoethyl}aminobenzoate

(1 ml). Dicyclohexylcarbodiimide (40 mg) was introduced.
 The material was stirred for 20h, in which time a solid

was precipitated. The material was filtered through a plug of glass wool, and the solvent removed by evaporation to leave a pale cream gum (288 mg). Assuming complete esterification, the product was about 45% pure by weight.

0155388

EXAMPLE 4

Active centre-linked conjugate of human tissue plasminogen activator with human serum albumin (LINKER EXAMPLE 1)

(a) N-ε-(4-thiobutyrimino)-[LYSINE] human serum albumin

Human serum albumin (Kabi, 200 mg/ml) was mixed with solid iminothiolane (Sigma, 10.3 mg) and incubated at pH 7.4, 37°C for 1h. The mixture was gel filtered into 3.5 ml 20% glycerol, 0.1 M Tris.HCl, 0.9% w/v NaCl buffer (TGS) pH 7.4 using a small Sephadex G-25 M column at 4°C. Active site titration using Ellmanns Reagent (1.0 mM, 20% TGS, pH 7.4) gave a thiol content of 2.33 mM (2 determinations), corresponding to an average of 1.09 thiols/mol HSA. The solution was stored at -40°C.

(b) N-ε-([2-[3-(4-butyriminodithio) propionyl] hydrazino] benzoic acid 4'-amidinophenyl ester.HCl)-[LYSINE] human serum albumin

Thiolated albumin (VI, 1.5 ml) was mixed with the acylating agent of Example I (b) (10 mM in DMSO, 0.5 ml, 1.43 molar excess) and incubated at 0°C for 60 min during which approximately 1.1 molar equivalents of pyridine 2-thione were released. Part of this material was subjected to two successive gel filtrations (Sephadex G-25 M) at 4°C into 1% w/v D-mannitol, 20 mM ammonium bicarbonate, 1.0 mM 6-aminohexanoic acid (MAE) buffer, pH 7.4. Final vol: 3.5 ml, 1.05 mM ester. The solution was stored at -40°C.

(c) Coupling to human tissue plasminogen activator (t-PA)

Purified t-PA was mixed with 0.1 M Tris.HCl pH 7.0 (0.5 ml) and the above

albumin ester (0.5 ml) added. On incubation at 0°C, amidolytic activity against S-2288 progressively decreased as illustrated in Figure 1. The acylation process appeared to have two phases: one relatively rapid ($t_1 \sim 13$ min) and the other slower ($t_2 \sim 98$ min). Additional inactivation occurred on freezing to -40°C and thawing. The product was mixed with a batch prepared under the same conditions but using ^{125}I -MPA.

(d) Purification of the conjugate

A column (2 x 55 cm) of Sephacryl S-300 was calibrated by chromatography of a mixture of unmodified albumin and ^{125}I -MPA (vial 5) in MAE buffer at 4°C and a flow rate of 22 ml/h. Albumin and ^{125}I eluted together at 62-88 ml. Crude conjugate chromatographed under the same conditions gave two peaks with apparent molecular weights greater than albumin and containing radioactivity. The fractions eluting at 42-64 ml were pooled and freeze-dried, yielding a white powder (244 mg). This process was repeated for several other batches.

(e) Deacylation of the conjugate

Freeze-dried batches of the conjugate were deacylated for pH 7.4, 37°C in the above TGS buffer containing 0.01% w/v Tween 80 detergent. The average first order deacylation rate constant for four batches was about $4 \times 10^{-4} \text{ sec}^{-1}$. On prolonged storage, a slower component of deacylation characterised by a rate constant of about $5 \times 10^{-5} \text{ sec}^{-1}$ became evident and this correlated with an increase in the average molecular weight of the preparation (see below).

Size of the conjugate

The size of the coupled product was examined by two techniques. First, polyacrylamide gel electrophoresis in the presence of sodium dodecyl sulphate (Laemmli, U.K. (1970), *Nature*, 227, 680) was used to separate the

components of the conjugate which were detected either by autoradiography of the ^{125}I -t-PA or by fibrin overlay zymography (Granelli-Piperno, A. & Reich, E. (1978), J. Exp. Med., 148, 223) which detects fibrinolytic enzymes. Secondly, size exclusion high performance liquid chromatography with calibrated modified silica columns was used to separate the conjugate mixture before and after deacylation. All these methods gave similar results which may be summarised as follows:

- (1) t-PA-associated radioactivity and enzymatic activity is associated with several species with a molecular weight in the range 130 to 900 kilodaltons. The individual components correspond to conjugates of t-PA with one, two, three etc (up to at least 10) albumin units.
- (2) All the discrete conjugate proteins appeared to possess fibrinolytic activity after deacylation. Treatment of the conjugate with TGS buffer or 0.5 M hydrazine pH 8.0 at 37°C for several hours resulted in the liberation of an active component (separable by HPLC or SDS-PAGE) with equimolar weight corresponding to unmodified t-PA (65-70 kilodaltons).
- (g) Clearance of the conjugate from the bloodstream of rats
(See also: Methods)

The rate of removal of unmodified t-PA and of albumin-t-PA conjugate from the circulation of rats was measured by both radiometric and fibrinolytic methods. The results are shown in Fig. 2. The activity of unmodified t-PA was

measured at 15 min). Radiometric

at 8 min, probably because of recycling of the ^{125}I -label into the bloodstream. Both the enzyme activity and the radioactivity of the albumin conjugate were released significantly more slowly than the unmodified

700

0155388

- 30 -

enzyme, with part of the dose being cleared with an apparent half-life of about 36 min. Integration of the concentration/time plots for the two agents suggests that the conjugate had at least a 10-fold greater bioavailability than the unmodified t-PA. Since the conjugate gives rise to unmodified t-PA by hydrolysis, it can therefore function as a slow-release form of the fibrinolytic agent in vivo.

EXAMPLE 5

- 31 -

Active centre-linked conjugate of human tissue-plasminogen
activator with human serum albumin (linker example 2)

Human serum albumin (Kabi, 200 mg/ml, 10 ml) was mixed with
2-iminothiolane (4.1 mg:1 mole equivalent) and incubated at
25°C for 75 min. Titration with Ellmann's reagent showed
a concentration of 2.82 mM thiol after this time. An
aliquot of this solution (0.5 ml) was mixed with the
acylating agent of Example 2(b) (70 µl of 20 mM in DMSO,
1 molar equivalent) and held on ice for 1h. t-PA (50,000
SU) and tracer ¹²⁵I-t-PA (960,000 c.p.m.) were added. The
mixture slowly decreased in activity over 72h at 0°C
although the decrease could be accelerated by incubation
at 25°C for short periods of time. The final amidolytic
activity was 37% of the initial level. The product was
applied to a column (300 x 22.5 mm) of TSK-G-3000 SW modified
silica gel and subjected to high performance gel permeation
chromatography. The buffer used was 0.08 M sodium phosphate,
0.32 M sodium chloride, 20% v/v ethanol pH 7.0 (Buffer H,
Hefti, F. (1982) Anal. Biochem., 121, 378-381) at a flow
rate of 1.0 ml/min at 22°C. Four peaks of protein-bound
radioactivity were observed, the first near the void volume
of the column (40 ml) was of very high molecular weight
and corresponded to MW around 150 kDa. The second peak at 52 ml was of material with

and corresponded to MW around 150 kDa. The third peak at 52 ml was of material with
t-PA eluted at about 74 ml. A pool of eluate from 52-66 ml

was collected, gel filtered into MAE buffer (diluted 1:5
v/v in water and containing 1 mg/ml human serum albumin)
and freeze-dried. This yielded 81 mg of a white solid.
Radiochemical yield: 16.5%. Deacylation of this material
5 in TGS buffer containing 0.01% w/v Tween 80, at 37°C
showed that it contained initially about 14% uncoupled
t-PA and that this rate constant for deacylation was about
 $6.4 \times 10^{-5} \text{ sec}^{-1}$.

This material was injected into rats at a dose of 2000 SU/
10 kg and the fibrinolytic activity in the bloodstream
determined as described in method (c). Fig. 3 shows
the clearance of the conjugate (2000 SU/kg) from the
bloodstream of rats (n = 5). A slow phase of
clearance with a half-life of about 36 min could be
15 identified. Comparison with Fig. 2 shows that this
clearance is appreciably slower than that of unmodified
t-PA.

EXAMPLE 6

- 33 -

Active-centre linked conjugate of human tissue plasminogen activator with human immunoglobulin G (linker example 1)

(a) N-ε-4-thiobutyrimino[lysine]human immunoglobulin G

5 Human immunoglobulin G (Sigma, 100 mg) was dissolved in degassed 50 mM sodium phosphate 0.1 M sodium chloride pH 7.4 buffer (PBS buffer, 1.0 ml) and 2-iminothiolane (50 µl of a fresh 50 mM solution in PBS) added. The mixture was incubated at 37°C for 1h and then gel filtered on a small
10 Sephadex G-25 column into 3.4 ml of PBS at 4°C. The product was used immediately.

(b) N-ε-(2-[3-(4-butyriminodithio)propionyl]hydrazino)benzoic acid 4'-amidinophenyl ester.HCl)-[lysine]-human immunoglobulin G

15 The above thiolated IgG was mixed with the acylating agent of Example 1(b) (85 µl of 20 mM in DMSO) and incubated at ambient temperature (22-24°C) for 10 min. Cold (-20°C) ethanol (10 ml) was added with mixing over 30 sec. The mixture was centrifuged at 18,000 g for 30 min at 4°C and
20 the pellet redissolved in PBS buffer (3.4 ml). Saturated ammonium sulphate solution (10 ml) was added with mixing
of the same buffer at 4°C. The product was used immediately.

(c) Coupling to human tissue plasminogen activator (t-PA)

Purified t-PA (approx. 26 nmoles) was mixed with the product of step (b) to give a solution containing 51,000 SU/ml. The mixture was held on ice for 24h after which the residual activity was 11,600 SU/ml (22.7%). The product was centrifuged to remove small amounts of insoluble material and an aliquot (50 μ l) diluted into TGS buffer containing 0.01% w/v Tween 80. The deacylation of this sample was measured at 37°C and gave a first order constant of about $5.7 \times 10^{-4} \text{ sec}^{-1}$. The remaining product was stored in solution at -70°C and showed a high molecular weight (c-200 Kilodaltons) fibrinolytic component (susceptible to hydrolysis) when analysed by fibrin overlay zymography.

EXAMPLE 7

- 35 -

Active-centre-linked conjugate of human tissue-plasminogen activator with human serum albumin (linker example 3)

(a) 4-{N-2-(3-[2-pyridyldithio]propionyl)aminoethylamino}-benzoyl human tissue-plasminogen activator

5 Tissue-plasminogen activator (46 nmoles in 3.0 ml 0.1 M sodium 4-guanidinobutyrate pH 7.4 was treated with the acylating agent of Example 3 (30 μ l of 50 mM in DMSO) at 4°C for 16h and then at 25°C for 5h. After this time, 4.6% of the initial amidolytic activity remained. 2.5 ml
10 of this solution was gel filtered into 3.5 ml of TGS buffer containing 0.01% w/v Tween 80. Dilution of an aliquot of this solution 1:20 v/v into TGS/Tween 80 and deacylation at 37°C indicated that the first order deacylation rate constant was about $6.6 \times 10^{-4} \text{ sec}^{-1}$.
15 The remaining solution was used immediately.

(b) Coupling to thiolated human serum albumin

Human serum albumin (Sigma, 200 mg) was dissolved in PBS (1.0 ml) and incubated for 1h at 25°C with 60 μ l of 50 mM fresh aqueous 2-iminothiolane. The product was gel
20 filtered into PBS (3.5 ml) and the filtrate immediately

SDS-PAGE/fibrin overlay zymography of the

product demonstrated the presence of high molecular weight fibrinolytic components which were susceptible to hydrolysis.

EXAMPLE 8

- 37 -

A conjugate of urokinase linked to the active centre of
human plasmin (linker example 1)

(a) Preparation of thiolated urokinase

5 High molecular weight urokinase (Serono 500,000 IU,
32.8 nmoles) was mixed with tracer ^{125}I -UK (1.13×10^6 c.p.m.)
in PBS buffer (1.0 ml) and treated with 2-iminothiolane
(25 μl of a 50 mM solution in cold (0°C) water). The mixture
was incubated at 25°C for 75 min and gel filtered (Sephadex^R
10 G-25) into MAE buffer diluted 1:1 v/v with water (3.5 ml).
This solution was freeze-dried.

(b) 4-(3-{2-pyridyldithio}propionyl)hydrazino benzoyl
(Ser-740) human plasmin

15 Lys-plasminogen (100 mg) was dissolved in TGS buffer (1.0 ml)
and activated with urokinase (5000 IU) at 0°C for 2h. The
resulting plasmin had an activity of 3.7×10^6 SU/ml and was
treated with the acylating agent of Example 1 (50 μl of a
20 mM solution in DMSO) for 1h at 0°C . After this time, the
amidolytic activity was reduced to 0.6% of the initial level
20 and the mixture was gel filtered into MAE buffer (3.4 ml)

and freeze-dried in 0.2 ml aliquots. One aliquot was

dithionitron (2 mM). The absorbance at 343 nm attributable to release of pyridine-2-thione indicated

that each aliquot contained approximately 27 nmoles of acyl-enzyme. The first order deacylation rate constant of the product was about $1.96 \times 10^{-4} \text{ sec}^{-1}$.

(c) Coupling of the components

5 The above urokinase was mixed with 5 aliquots of acyl-enzyme in PBS buffer (1.0 ml) and held at 0°C for 2h. Approximately 61 nmoles of pyridine 2-thione were released as measured spectrophotometrically. The product was purified by preparative high performance gel permeation chromatography
10 as described in Example 5. The conjugate eluted at around 60-68 ml and this pool was gel filtered into MAE buffer (diluted 1:1 v/v with water and containing 1 mg/ml human serum albumin and freeze-dried in 6 aliquots). Unmodified urokinase eluted from the column at around 78 ml. The final
15 yield of conjugate was 8.3% (radiochemical) or 9.6% (amidolytic). Hydrolysis of the conjugate at 37°C for 45 min in TGS buffer indicated release of plasmin as measured by a 4-fold increase in amidolytic activity against the plasmin substrate S-2251. SDS PAGE and fibrin overlay zymography
20 revealed components corresponding to one and two molecules of acyl plasmin conjugated to each UK molecule. The high MW bands were sensitive to hydrolysis in TGS buffer at 37°C for 3h.

EXAMPLE 9

A conjugate of p-aminobenzoyl urokinase linked to the active centre of human plasmin (linker example 2)

Urokinase (Serono 500,000 IU, 32.8 nmoles) was mixed with
5 tracer ^{125}I -UK (1.70×10^6 c.p.m.) and treated with 2-
iminothiolane (13.3 μl of 10 mM in cold water) in PBS buffer
(0.5 ml) at 25°C for 75 min. This product was used without
further processing. 2-(N-6-[3-{2-pyridyldithio}propionyl]
aminoethylaminobenzoyl (Ser-740) human plasmin was prepared
10 by reacting plasmin (prepared as in Example 8(b), 0.9 ml)
with the acylating agent of Example 2(b) (80 μl of 20 mM
in DMSO) for 2h at 25°C . The amidolytic activity of the plasmin
decreased to 8.8% of the initial activity under these
conditions. The product was gel filtered into MAE buffer,
15 diluted 1:5 v/v in water (3.2 ml) and freeze-dried in 4
aliquots. Reduction of one of these aliquots with dithio-
threitol indicated an acyl-enzyme content of about 140 nmoles.
The above thiolated urokinase was mixed with 4-aminobenzoic
acid 4'-amidinophenyl ester.HCl (20 μl of 50 mM solution in
20 DMSO) and added to 1 vial of acyl-plasmin. The mixture was
incubated at 25°C for 40 min and then chromatographed using
the HPLC conditions described in Example 4. Monitoring the
radioactivity of the effluent showed that the conjugate
eluted between 59 and 66 ml whereas unmodified urokinase

freeze-dried in aliquots. Deacylation of one of these

aliquots in TGS buffer at 37°C indicated a first order
deacylation rate constant for removal of the p-aminobenzoyl
group at the urokinase active site of about 2.7×10^{-4}
sec⁻¹. The acyl group linking the urokinase to the plasmin
was much more stable ($k < 2 \times 10^{-5}$ sec⁻¹). SDS-PAGE and
fibrin overlay zymography showed a pattern similar to that
noted for Example 8.

The compound was injected into 4 rats (at 8,000 SU/kg i.v.)
and the clearance of fibrinolytic activity from the
bloodstream measured using the technique described in
Method (c) and compared with the clearance of unmodified
urokinase (at 40,000 SU/kg). Figure 4 shows the results
obtained using euglobulin precipitation. An extended
plasma half-life of 33 - 47 min compared with
unmodified UK was observed. The clearance of
this agent was also studied in guinea pigs (Method (e) at
a dose of 5,000 SU/kg i.v. Fig. 5 compares the clearance
of unmodified UK and the conjugate and shows that a half-
life of 40-50 min applies to at least 50% of the injected
conjugate dose in marked contrast to the more rapid
clearance of UK itself.

EXAMPLE 10

A conjugate of urokinase linked to the active site of
human plasmin (linker example 3)

(a) 4-{N-2-(3-[2-pyridyldithio]propionyl)aminoethylamino}
5 benzoyl human plasmin (method 1)

Human plasmin solution was prepared by the method of
Example 8(b) and had an amidolytic activity of 3.05×10^6
SU/ml. 1.5 ml of this solution was mixed with the
acylating agent of Example 3 (f) (75 μ l of a 50 mM solution in
10 DMSO). The mixture was incubated at 4°C for 16h, at 25°C
for 2h and, after a further 75 μ l of acylating agent had
been added, at 37°C for 30 min. The amidolytic activity
was then reduced to 1.16% of the initial level. The
product was filtered through a 0.22 μ filter and purified
15 by HPLC using the conditions described for Example 5. The
acyl plasmin eluted between 62 and 68 ml and was well
separated from excess acylating agent. The acyl-plasmin
pool was gel filtered into MAE buffer (9.6 ml) and freeze-
dried in 6 aliquots. Reduction of one such aliquot with
20 dithiothreitol indicated an acyl-enzyme content of 5.62
nmoles/aliquot. Deacylation of the acyl-enzyme in TGS/
Tween 80 buffer at 37°C gave a first order deacylation rate
constant of about $6.4 \times 10^{-5} \text{ sec}^{-1}$.

(b) Coupling to thiolated urokinase

High molecular weight urokinase (Serono, 500,000 IU, 32.8 nmoles) was mixed with tracer ^{125}I -UK (1.33×10^6 c.p.m.) in PBS buffer containing 0.01% w/v Tween 80 (1.0 ml) and mixed with 2-iminothiolane (50 μl of 50 mM fresh in cold water), followed by incubation at 25°C for 1h. The product was gel filtered into PBS/Tween 80 (3.5 ml) and immediately added to 5 vials of the above acyl-plasmin (28.1 nmoles) and 0.2 ml of a 44 μM solution of bovine lung trypsin inhibitor (aprotinin). The mixture was placed in a CentriconTM centrifugal ultrafiltration vial and centrifuged at 4°C for 2h at 5000 g. This process reduced the volume to 1.1 ml. The product was purified by HPLC using the conditions described for Example 4. Three peaks of radioactivity were observed: the first at about 56ml corresponded to a conjugate of MW \sim 200 kilodaltons, the second at about 64 ml to material of 100-140 kilodaltons and the third at 80-82 ml to unmodified urokinase. The first two peaks (pools: 52-58 ml, 60-68 ml) were gel filtered into MAE buffer and freeze-dried. The radiochemical yield of the high MW fraction (corresponding to urokinase coupled to 2 plasmin molecules) was 6.3% and that of the second fraction (a 1:1 conjugate) was 9.8%. The recovery of amidolytic activity in the two fractions was 3.9 and 11% respectively. SDS-PAGE/fibrin overlay zymography confirmed the

EXAMPLE 11

- 43 -

A conjugate of human tissue-plasminogen activator linked to the active centre of human plasmin (linker example 1)

Tissue-plasminogen activator (18,000 SU) mixed with tracer ^{125}I t-PA (2×10^6 c.p.m.) in PBS buffer containing 0.01% w/v Tween 80 (2.5 ml) was treated with 2-iminothiolane (25 μl of 50 mM in cold PBS/Tween) and the mixture incubated at 25°C for 30 min. The product was gel filtered into PBS/Tween buffer (3.0 ml) and mixed with one aliquot of the acyl-plasmin of Example 8(b) and aprotinin solution (20 μl of 44 μM). The mixture was incubated on ice for 1h. SDS-PAGE/fibrin overlay zymography of this material revealed a high MW conjugate band which was sensitive to hydrolysis.

EXAMPLE 12Active centre linked conjugate of tissue-plasminogen
activator with human plasminogen (linker example 1)

Human lys-plasminogen (1.2 ml of a 9.4 mg/ml solution in PBS buffer containing 33% v/v glycerol) was mixed with 2-iminothiolane (24 μ l of a fresh 50 mM solution in water) and incubated at 37°C for 4 min. The acylating agent of Example 1(b) (150 μ l of a 10 mM solution DMSO) was added and the mixture held on ice for 1h. The product was gel filtered (SephadexTM G-25) into PBS buffer containing 1.0 mM 6-aminohexanoic acid (3.2 ml). Part of the resulting thiolated plasminogen solution (1.0 ml) was mixed with t-PA (20,000) and tracer ¹²⁵I-t-PA (1 x 10⁶ c.p.m.) and incubated at 0°C. After 2½h, the amidolytic activity of the t-PA was decreased to 7.7% of the initial level. Part of the product was subjected to HPLC separation on a modified silica gel column (TSK G-3000, 600 x 7.5 mm) in Buffer H (see Example 5) at 0.5 ml/min and 22°C. A high molecular weight peak of radioactivity eluted at about 17 ml and unmodified t-PA eluted at about 20 ml. SDS-PAGE/fibrin zymography also revealed the presence of a hydrolysis-sensitive high MW fibrinolytic component. (

EXAMPLE 13

- 45 -

A conjugate of human tissue-plasminogen activator linked to the active centre of human plasmin (linker example 3)

Tissue-plasminogen activator (78 nmoles in 3.4 ml 0.1 M sodium 4-guanidinobutyrate, 0.3 M sodium chloride, 20 mM sodium phosphate, 0.01% w/v Tween 80 pH 7.4) was treated with 50 μ l of 50 mM 2-iminothiolane (fresh in cold water) at 25°C for 70 min. 1.7 ml of this solution was gel filtered into 3.4 ml of the above buffer on SephadexTM G-25. Titration of the thiol groups in the t-PA with Ellmann's reagent indicated an average of about 0.6-0.7 moles of thiol/mole of protein. The thiolated t-PA was mixed with aprotinin (132 nmoles), the above buffer (0.4 ml) and the acyl-plasmin of Example 8(b) (53 nmoles). The mixture was concentrated in CentriconTM vials to a final volume of 1.0 ml (5000 g/4°C for 2h). The product contained about 23 nmoles of t-PA as measured by amidolytic assay. Analysis of the product by SDS-PAGE/fibrin overlay zymography showed that a fibrinolytically active component with an apparent molecular weight of about 150 kilodaltons was present together with unmodified t-PA. Reduction of this material with dithiothreitol (c. 20 mM at 0°C for 30 min) removed most of the high molecular weight component, thus confirming that the components of the conjugate were linked

EXAMPLE 14

- 46 -

A conjugate of urokinase linked to the active centre of human plasmin (linker example 2, no acylation of the active centre of urokinase)

5 High MW urokinase (Serono, 500,000 IU, 32.8 nmoles) was dissolved in PBS buffer containing 0.01% w/v Tween 80 (1.0 ml) and treated with 25 μ l of 50 mM 2-iminothiolane at 25°C for 1h. The product was gel filtered into MAE buffer and freeze-dried. Ellmann's titration showed that
10 the average thiol content of the modified UK was 0.8 moles/mole. The product (64 nmoles) was mixed with N-2-[N-6-{3-(2-pyridyldithio)propionyl}aminohexyl]aminobenzoyl (SEK 740) human plasmin prepared as in Example 9 (330 nmoles) in PBS/Tween buffer (1.0 ml) and held at 0°C for 4h. The
15 product was partially purified by HPLC on a modified silica gel column (TSK G-4000 SW, 7.5 x 600 mm) in buffer H at a flow rate of 1.0 ml/min at 25°C. The product eluting between 20 and 24 ml was gel filtered into MAE buffer containing 1 mg/ml human serum albumin and freeze-
20 dried. 151 g of white solid resulted and part of this (81 mg) was repurified on a column (16 x 200 mm) of Sephacryl TM S-300 in 4 mM ammonium bicarbonate, 0.1% w/v D-mannitol at a flow rate of 10 ml/h at 4°C. Two fractions were collected and freeze-dried. The first corresponded

SDS-PAGE/fibrin overlay zymography showed that the high MW

fraction contained a fibrinolytically active component at
around 220 kilodaltons (UK linked to 2 plasmin molecules)
as well as the conjugate of UK with one plasmin
(134 kilodaltons) and free UK. Like the conjugate of
Example 9, the product was not easily hydrolysed.

Claims


- 1) A derivative of a fibrinolytic enzyme in which the catalytic site on the enzyme which is responsible for fibrinolytic activity is blocked by a human protein attached thereto by way of a reversible linking group.
- 2) A derivative according to claim 1 in which the catalytic site on the enzyme is blocked by a group of formula (I):



in which



P is a human protein modified by treatment with an amino acid side chain specific reagent to include a protein attachment group;

X is an acyl group of formula



wherein R is an aliphatic or aromatic residue,
A is a bridging group comprising at least one heteroatom selected from oxygen sulphur and nitrogen, in which the nitrogen is optionally substituted by C₁-6 alkyl,
and B is a hydrophilic linking group, linked to the

3) A derivative according to claim 2 wherein group B is a



4) A derivative according to any one of the preceding claims wherein the fibrinolytic enzyme is tissue type plasminogen activator, urokinase or plasmin.

5) A derivative according to any one of the preceding claims wherein the human protein is human serum albumin, immunoglobulin, human fibrinogen, human plasminogen or a fibrinolytic enzyme or proenzyme thereof.

6) A derivative according to claim 1 which is

human serum albumin \rightarrow human tissue plasminogen
activator

human serum albumin \rightarrow urokinase

urokinase or acylated urokinase \rightarrow human plasmin

human tissue plasminogen activator \rightarrow human plasmin

human immunoglobulin G \rightarrow human tissue plasminogen
activator


where \rightarrow represents a reversible linkage to the active centre of the enzyme.

7) A process for preparing a derivative according to any one of the preceding claims which process comprises reacting together a human protein optionally modified to include a protein attachment group, a fibrinolytic enzyme and a linking agent having a moiety capable of reacting with the catalytic site of the enzyme and a moiety capable of reacting with a protein amino group or protein attachment group to form a reversible linking group.

8) A pharmaceutical composition comprising a derivative of a fibrinolytic enzyme according to claim 1 in combination with a pharmaceutically acceptable carrier.

9) A derivative of fibrinolytic enzyme as defined in claim 1 for use in the treatment of thrombotic diseases.

10) A compound of formula II:



in which B, A and X are as defined in claim 2, and W represents a group capable of reacting directly with the amino acid side chain of a protein or protein attachment group, Z is a counter anion and R₁ to R₄ are each independently hydrogen or an electron withdrawing

1/4




Fig. 1




Fig. 2





Fig. 3

3/4

▲ Urokinase (40,000 SU/kg, n=8)

• Example 9 (5000 SU/kg, n=3)






Fig 5